

a new approach to Goldbach conjecture and goldbach numbers density

Nasroallah Hitar

January 2023

Abstract

in this paper we will provide a simple prove of the existence of infinitely many Goldbach numbers.

1 Introduction

The Goldbach Conjecture is a famous unsolved problem in number theory that states that every even integer greater than 2 can be expressed as the sum of two prime numbers. It was first proposed by the German mathematician Christian Goldbach in a letter to the Swiss mathematician Leonhard Euler in 1742. Despite much effort, a proof or counterexample has yet to be found. It is one of the oldest unsolved problems in number theory and in all of mathematics.

2 Demonstration principle

Let

$$P_G(n) : (\exists p \in \mathbb{P} : 2n - p \in \mathbb{P})$$

$$G := \{n \in \mathbb{N} | P_G(n)\}$$

let's suppose that G is finite.

We have that $2 \in G$.

$$\because 2 \cdot 2 - 2 \in \mathbb{P}$$

and we have that $G \subset \mathbb{N}$, and (G is finite).

Peano axiome : G has an element maximal, let's note it as g , $\max(G) := g$.

let $p \in \mathbb{P} : p > 2, \forall m \in \mathbb{N} : m > g : 2m = p + q$
such that $q \notin \mathbb{P}$ and $q \notin 2\mathbb{N}$ ($\because p$ is an odd prime.)

let $k \in \mathbb{N} : 2m + 2k = p + q + 2k$
 $\implies 2(m + k) = p + (q + 2k)$, and we have that :
 $2(m + k) \geq 2m > 2g \implies 2(m + k) \notin G$

$$\therefore q + 2k \notin \mathbb{P} \tag{1}$$

(because if we suppose that :
 $q + 2k \in \mathbb{P} \implies 2(m + k) - p \in \mathbb{P} \therefore 2(m + k) \in G$
and that's impossible. because g is the maximal element
of G . and $m > g$.)

let

$$f_q : \mathbb{N} \longrightarrow 2 \llbracket q, +\infty \llbracket + 1$$

$$q \longrightarrow q + 2k$$

we have that $q + 2k = q + 2k' \implies k = k'$

so f_q : is a injective map.

let : $\lambda \in 2 \llbracket q, +\infty \llbracket + 1$ so $\exists k' \in \llbracket q, +\infty \llbracket : \lambda = 2k' + 1$ to
prove that f_q is a surjective map, we have to prove the
existence of $k \in \mathbb{N}$ such that :

$$f_q(k) = 2k' + 1 \implies q + 2k = 2k' + 1$$

$\implies k = \frac{2k'+1-q}{2} \in \mathbb{N} (\because q \text{ is an odd integer } .)$

the map $f_q : \mathbb{N} \rightarrow \mathbb{N}$ is a bijective map in

$2 \llbracket q, +\infty[+ 1$. using equation (1) we have that

$$\forall k \in \mathbb{N} : q + 2k \notin \mathbb{P}$$

in other way :

$\forall k \in \mathbb{N} : f_q(k) \notin \mathbb{N} \implies \forall \lambda \in 2 \llbracket q, +\infty[+ 1 : \lambda \notin \mathbb{P}$

hence ! contradiction . because there exist infinitely many primes.

the set G is infinite set . QED.